

**ATLAS OF TUMOR PATHOLOGY**

Third Series  
Fascicle 13

**TUMORS OF THE  
LOWER RESPIRATORY TRACT**

by

**THOMAS V. COLBY, M.D.**

Laboratory of Medicine and Pathology  
Mayo Clinic Scottsdale  
Scottsdale, Arizona 85259

**MICHAEL N. KOSS, M.D.**

Co-Chairman  
Department of Pulmonary and Mediastinal Pathology  
Armed Forces Institute of Pathology  
Washington, D.C. 20306-6000  
and  
Department of Pathology  
University of Southern California  
Los Angeles, California 90033

**WILLIAM D. TRAVIS, M.D.**

Co-Chairman  
Department of Pulmonary and Mediastinal Pathology  
Armed Forces Institute of Pathology  
Washington, D.C. 20306-6000

Published by the  
**ARMED FORCES INSTITUTE OF PATHOLOGY**  
Washington, D.C.

Under the Auspices of  
**UNIVERSITIES ASSOCIATED FOR RESEARCH AND EDUCATION IN PATHOLOGY, INC.**  
Bethesda, Maryland  
1995

Accepted for Publication  
1994

---

Available from the American Registry of Pathology  
Armed Forces Institute of Pathology  
Washington, D.C. 20306-6000  
ISSN 0160-6344  
ISBN 1-881041-17-4

2063632651

# TUMORS OF THE LOWER RESPIRATORY TRACT

## Contents

1. Introduction and Classification .....	1
WHO Histologic Typing of Lung Tumors .....	1
2. Embryology, Anatomy, and Congenital, Developmental, and Related Lesions .....	3
Embryology .....	3
Anatomy and Histology .....	3
Congenital Anomalies and Related Lesions .....	11
Bronchial Atresia (Regional or Segmental Pulmonary Overinflation) .....	12
Pulmonary Sequestration (Intralobar Sequestration, Extralobar Sequestration) .....	12
Congenital Bronchopulmonary Foregut Malformations .....	16
Bronchogenic Cysts (Bronchial Cysts) .....	16
Congenital Cystic Adenomatoid Malformation .....	20
Infantile Lobar Emphysema/Polyalveolar Lobe .....	25
Pulmonary Interstitial Emphysema .....	25
Diffuse Pulmonary Lymphangiectasia .....	25
Ectopic Tissues in the Lung .....	25
3. Specimen Handling and Special Techniques .....	31
Routine Handling of Lung Specimens .....	31
The Surgical Pathology Report .....	32
Special Techniques .....	33
Immunohistochemistry .....	33
Flow Cytometry .....	34
Molecular Biology .....	35
4. Papillary Tumors of the Bronchial Tree .....	49
Squamous Papilloma and Papillomatosis .....	49
Other Bronchial Papillomas .....	55
Columnar Papillomas .....	55
Inverted Papillomas or Polyps .....	55
Transitional Cell Papillomas .....	55
Fibroepithelial Polyps .....	55
5. Miscellaneous Benign Epithelial Tumors .....	57
Papillary Adenoma of Type 2 Cells .....	57
Alveolar Adenoma .....	59
Mucinous Cystadenoma and Pulmonary Mucinous Tumors of Borderline Malignancy ..	61
Mucinous Cystadenoma .....	61
Mucinous Cystic Tumor of Borderline Malignancy .....	61
6. Tumors of Salivary Gland Type .....	65
Mucoepidermoid Carcinoma of the Bronchus .....	65
Adenoid Cystic Carcinoma .....	72
Mucous Gland Adenoma .....	79
Pleomorphic Adenoma .....	82
Acinic Cell Carcinoma .....	84
Other Rare Salivary Gland Tumors .....	85
Carcinoma ex Pleomorphic Adenoma (Malignant Mixed Tumor) .....	85

2063632652

Pulmonary Oncocytoma .....	86
Myoepithelioma .....	86
Adenosquamous Carcinoma with Amyloid-like Stroma .....	87
7. Carcinoma of the Lung: Overview, Incidence, Etiology, and Screening .....	91
Overview .....	91
Incidence .....	93
Etiology and Pathogenesis .....	93
Screening for Carcinoma of the Lung .....	97
8. Carcinoma of the Lung: Clinical and Radiographic Aspects, Spread, Staging, Management, and Prognosis .....	107
Demographics .....	107
Signs and Symptoms .....	107
Radiographic Features .....	110
Patterns of Spread .....	112
Staging, Management, and Prognosis .....	117
9. Morphologic Diagnosis and Heterogeneity of Carcinoma of the Lung .....	135
Diagnosis .....	135
Histologic Heterogeneity .....	136
10. In Situ and Early Invasive (Occult) Squamous Cell Carcinoma .....	145
11. Squamous Cell Carcinoma and Variants .....	157
12. Adenocarcinoma of the Lung (Excluding Bronchioloalveolar Carcinoma) .....	179
13. Bronchioloalveolar Carcinoma .....	203
14. Small Cell Carcinoma and Large Cell Neuroendocrine Carcinoma .....	235
Small Cell Carcinoma .....	235
Large Cell Neuroendocrine Carcinoma .....	248
15. Large Cell Carcinoma .....	259
16. Adenosquamous Carcinoma, Carcinomas Associated with Cysts, and Paget Disease of the Bronchus .....	279
Adenosquamous Carcinoma .....	279
Carcinomas Associated with Lung Cysts .....	282
Pagetoid Spread of Adenocarcinoma in the Bronchus .....	285
17. Carcinoid and Other Neuroendocrine Tumors .....	287
Unifying Concept of Pulmonary Neuroendocrine Tumors .....	287
Neuroendocrine Cells .....	287
Classification .....	287
Neuroendocrine Body .....	289
Carcinoid Tumorlet .....	290
Typical and Atypical Carcinoid Tumors .....	294
Unusual Lung Tumors with Neuroendocrine Differentiation .....	309
Paranglioma .....	309
Primitive Neuroectodermal Tumor (Malignant Small Round Cell Tumor of the Thoracopulmonary Region) .....	310
Neuroendocrine Carcinoma with Rhabdoid Phenotype .....	311
Amphicrine Neoplasms .....	311
Neuroendocrine Carcinoma with Anemone Features .....	313
Pulmonary Blastoma with Neuroendocrine Differentiation .....	313
18. Hamartoma .....	319

2063632653

19. Fibrous and Fibrohistiocytic Tumors and Tumor-Like Conditions	327
Inflammatory Pseudotumor (Plasma Cell Granuloma-Histiocytoma Complex)	327
Intrapulmonary Localized Fibrous Tumor	338
Malignant Fibrous Histiocytoma	342
Miscellaneous Fibrous and Fibrohistiocytic Tumors	345
Fibrosarcomas	345
Mesenchymal Cystic Hamartoma	348
Cystic Fibrohistiocytic Tumor	348
20. Miscellaneous Mesenchymal Tumors	353
Smooth Muscle Tumors	353
Leiomyoma and Leiomyosarcoma	353
Lymphangioliomyomatosis	356
Glomus Tumors	360
Skeletal Muscle Tumors	361
Rhabdomyoma	361
Rhabdomyosarcoma	361
Vascular Tumors and Related Conditions	361
Vascular Malformations	361
Hemangioma/Hemangiomatosis	361
Lymphatic Lesions	363
Epithelioid Hemangioendothelioma	363
Kaposi Sarcoma	368
Pulmonary Angiosarcoma	371
Pulmonary Artery and Vein Sarcomas	371
Hemangiopericytoma	378
Neurogenic Tumors	378
Neurilemmoma and Neurofibroma	378
Neuroma and Ganglioneuroma	378
Malignant Nerve Sheath Tumors	378
Malignant Psammomatous Melanotic Schwannoma	380
Neuroblastoma and Ganglioneuroblastoma	380
Meningioma	380
Cartilaginous Tumors of the Lung	382
Chondroma	382
Chondroblastoma	384
Chondrosarcoma	384
Osteogenic Tumors	385
Metaplastic Bone	385
Osteosarcoma	385
Fatty Tumors	386
Lipomas	386
Liposarcomas	388
Aveolar Soft Part Sarcoma	388
21. Mixed Epithelial and Mesenchymal Tumors	395
Pulmonary Blastoma	395
Well-Differentiated Fetal Adenocarcinoma (Pulmonary Endodermal Tumor)	395
Biphasic Pulmonary Blastoma	403

2063632654

Cystic and Pleuropulmonary Blastomas of Childhood .....	406
Carcinosarcomas of the Lung .....	411
22. Lymphoreticular Disorders .....	419
Histologic Diagnosis of Lymphoreticular Infiltrates in the Lung .....	419
Lymphatics and Lymphoid Tissue of the Lung .....	420
Benign/Hyperplastic Disorders .....	422
Pseudolymphoma (Nodular Lymphoid Hyperplasia) .....	422
Lymphocytic Interstitial Pneumonia and Diffuse Lymphoid Hyperplasia .....	424
Giant Lymph Node Hyperplasia (Castleman Disease) .....	427
Angioimmunoblastic Lymphadenopathy .....	427
Lymphomas Presenting in Lung .....	428
Small Lymphocytic and Lymphoplasmacytoid Lymphomas (Including Low-Grade Lymphomas of BALT) .....	428
Angiocentric Immunoproliferative Lesions/Angiocentric Lymphomas (Lymphomatoid Granulomatosis/Polymorphic Reticulosis) .....	434
Large Cell Lymphoma of Lung .....	443
Intravascular Lymphomatosis (Angiotropic Lymphoma) .....	443
Other Non-Hodgkin Lymphomas of the Lung .....	445
Primary Pulmonary Hodgkin Disease .....	445
Plasmacytoma .....	449
Mast Cell Tumor .....	449
Systemic Lymphoproliferative Disorders Secondarily Involving the Lung .....	450
Hodgkin Disease and Non-Hodgkin Lymphoma .....	450
Cutaneous T-Cell Lymphomas (Mycosis Fungoides and Sézary Syndrome) .....	450
Malignant Histiocytosis .....	451
Multiple Myeloma .....	451
Systemic Light Chain Disease .....	453
Waldenström Macroglobulinemia .....	453
Post-Transplant Lymphoproliferative Disorders and Related Conditions .....	454
Leukemic Infiltrates Involving the Lung .....	454
Myelofibrosis .....	459
Pleural Involvement in Lymphoreticular Disorders .....	459
Pulmonary Histiocytosis X .....	459
23. Miscellaneous Tumors and Tumors of Uncertain Histogenesis .....	465
Sclerosing Hemangioma of the Lung .....	465
Clear Cell Tumor .....	471
Minute Pulmonary Meningothelial-Like Nodules ("Minute Pulmonary Chemodectomas") .....	474
Granular Cell Tumor .....	478
Thymoma .....	482
Malignant Melanoma .....	483
Germ Cell Tumors .....	487
Teratoma .....	488
Choriocarcinoma .....	490
Ependymoma .....	490
24. Tumor-Like Conditions .....	495
Nodular Pulmonary Amyloidosis .....	495
Pulmonary Hyalinizing Granuloma .....	501

2063632655

## Contents

Pulmonary Malakoplakia .....	506
Rounded Atelectasis .....	510
Focal Organizing Pneumonia .....	513
25. Tumors Metastatic to the Lung .....	517
Patterns of Metastasis .....	517
Multiple Nodules .....	517
Lymphangitic Metastasis .....	517
Endobronchial Metastasis .....	524
Metastatic Tumor Embolization .....	526
The Solitary Pulmonary Metastasis .....	530
Pleural Metastases .....	531
Distinctive Histologic Growth Patterns .....	533
Lepidic Growth Pattern .....	533
Interstitial Spread .....	534
Cavitation .....	534
Special Techniques and Their Application in the Diagnosis of Pulmonary Metastases .....	538
Immunohistochemistry .....	538
Electron Microscopy .....	541
Oncogenes .....	541
Index .....	547

2063632656

# CARCINOMA OF THE LUNG: OVERVIEW, INCIDENCE, ETIOLOGY, AND SCREENING

## OVERVIEW

The common tumors generally included under the heading of carcinoma of the lung are squamous cell carcinoma, adenocarcinoma, small cell carcinoma, and large cell carcinoma. These tumors share many clinical, etiologic, and demographic features. Bronchioloalveolar carcinoma is included as a subset of adenocarcinoma, although it has distinctive features of its own.

The incidence of the major subtypes of lung carcinoma from several large studies is shown in Table 7-1. Most recent series suggest that adenocarcinoma is now more frequent than squamous cell carcinoma (fig. 7-1) (4,12); in fact, adenocarcinoma represented 56 percent of the lung carcinomas seen at Johns Hopkins Hospital from 1984 to 1987 (7). Adenocarcinoma is the most frequent subtype seen in women. A relative decrease in the incidence of squamous cell carcinoma has accompanied the increased incidence of adenocarcinoma from the 1960s through the 1980s (1,3,6, 12,13). Although not as dramatic as the change

Table 7-1

## FREQUENCY OF SUBTYPES OF LUNG CARCINOMA

Cell Type	Percent of Cases
A. Series prior to 1985 (21,139 cases)*	
Squamous cell carcinoma	37.7
Adenocarcinoma	24.9
(Bronchioloalveolar carcinoma: 2-2.8%)**	
Small cell carcinoma	18.7
Large cell carcinoma	18.7
B. SEER data 1983-1987 (59,260 cases)†	
Squamous cell carcinoma	30.0
Adenocarcinoma	32.2
(Bronchioloalveolar carcinoma: 0.7%)	
Small cell carcinoma	18.2
Large cell carcinoma	9.7
Other/unspecified	9.9

\*From references 3,6,11,13 primarily reflecting data from prior to 1985.

\*\*From references 10,13.

†From reference 12.




Figure 7-1

## INCIDENCE OF LUNG CARCINOMA

The changing incidence of squamous cell carcinoma, adenocarcinoma, bronchioloalveolar carcinoma, and large cell carcinoma is shown for three time periods between 1962 and 1986. The period from 1983 to 1986 is represented by 100 consecutive cases seen by the National Cancer Institute, whereas data for the two earlier periods are from the literature. (Fig. 1 from Gazdar AF, Linnoila RI. The pathology of lung cancer—changing concepts and newer diagnostic techniques. *Semin Oncol* 1988;15:215-25.)

2063632657

in the incidence of adenocarcinoma, the incidence of small cell carcinoma has also increased in recent years (12).

A relative decrease in the percent of central tumors compared to peripheral tumors has accompanied the change in histologic subtypes. Auerbach and Garfinkel (1) found a decrease in central tumors from 69.3 to 57.3 percent when comparing tumors seen prior to 1978 to those seen between 1986 and 1989.

The change in incidence in lung carcinoma subtypes may, in part, be related to changing histologic criteria in addition to a real change in incidence (4). Another complicating factor is the histologic heterogeneity of lung carcinomas, discussed in chapter 9.

Multiple separate primary carcinomas of the lung are well recognized; the incidence ranges from 0.2 to 2.0 percent of patients with lung carcinoma (2,9). Synchronous lung carcinomas are seen in about 2 percent of surgical resections (2). Of 50 cases reported by Martini and Melamed (8), 18 were synchronous and 32 were metachronous, with the time to diagnosis of the second tumor varying from 4 months to 16 years. The criteria for the diagnosis of two (or more) separate primary lung carcinomas used by Martini and Melamed are shown in Table 7-2. Squamous cell carcinoma is the most common subtype associated with multiple primary carcinomas of the lung, but many combinations occur (Table 7-3).

While gross and histologic features can be used to recognize synchronous primary lung carcinomas, other techniques may also be useful. Ichinose et al. (5) have shown that DNA flow cytometric patterns can separate synchronous primaries from intrapulmonary metastases.

Table 7-2

**CRITERIA FOR DIAGNOSIS OF  
MULTIPLE PRIMARY  
LUNG CARCINOMAS\***

**Metachronous tumors**

1. Histologically different\*\*
2. Similar histology, but
  - a. free interval between tumors of at least 2 years
  - b. origin from carcinoma in situ
  - c. second tumor in different lobe, but
    - (1) no carcinoma in lymphatics common to both tumors
    - (2) no extrapulmonary metastases at time of diagnosis

**Synchronous tumors**

1. Tumors physically distinct and separate
2. Histology
  - a. different\*\*
  - b. same, but in different segments if
    - (1) origin from carcinoma in situ
    - (2) no carcinoma in lymphatics common to both
    - (3) no extrapulmonary metastases at time of diagnosis

\*Modified from reference 8.

\*\*Adequate sampling mandatory.

Table 7-3

**HISTOLOGY OF MULTIPLE PRIMARY LUNG CARCINOMAS\***

Cell Type		Synchronous	Metachronous	Total
Tumor #1	Tumor #2			
Squamous	Squamous	31	59	90
Squamous	Adenocarcinoma	21	10	31
Squamous	Small cell	22	13	35
Squamous	Large cell**	5	6	11
Adenocarcinoma	Adenocarcinoma	4	9	13
Adenocarcinoma	Small cell	5	1	6
Adenocarcinoma	Large cell**	1	2	3
Small cell	Large cell**	1	0	1
Large cell**	Large cell**	4	1	5
		94	101	195

\*Modified from reference 8 (195 cases).

\*\*Includes some unspecified "anaplastic carcinomas."



## INCIDENCE

Data from the American Cancer Society show that in 1989 carcinoma of the lung was the most common cause of cancer death in both men (34.1 percent) and women (19.9 percent) (14). It was estimated that 170,000 new cases of lung carcinoma would occur in the United States in 1993. Based on data from the United States for the year 1989, 23.1 percent of all deaths were due to cancer, and 27.6 percent of the cancer deaths were due to carcinoma of the lung. Thus, approximately 6.4 percent of all deaths in the United States in 1989 were due to carcinoma of the lung. The age-adjusted cancer death rate for 1989 was close to 30 per 100,000 for women and 75 per 100,000 for men.

Death rates from lung carcinoma have risen steadily in both men and women since 1930, with the rise occurring earlier and being steeper in men. However, since 1970 there has been a sharp increase in the death rate in women (14). Evidence suggests that the rate may be leveling off or decreasing in men as a result of public health efforts to curb cigarette smoking (15).

## ETIOLOGY AND PATHOGENESIS

Cigarette smoking is the major cause of lung carcinoma in the United States and around the world (38). The lung carcinoma rate parallels smoking prevalence. There is a dose-response association between the number of cigarettes smoked and the risk of carcinoma of the lung, although changes in the composition of cigarettes and the introduction of filter tips in the 1950s and 1960s have lowered the risk of lung carcinoma 20 to 50 percent compared to earlier "high yield" cigarettes (30). The risk of lung carcinoma is also increased in individuals who are pipe or cigar smokers. The increased risk in smokers is seen for all types of lung carcinoma and decreases exponentially over time after cessation of cigarette smoking (38).

There are some variations in the histologic subtypes among smokers and nonsmokers (Table 7-4), with squamous cell carcinoma and small cell carcinoma showing the highest association with smoking. It is possible that the recent decrease in the frequency of squamous cell carcinoma is in part attributable to changing smoking habits (1).

Table 7-4

## HISTOLOGIC SUBTYPES OF CARCINOMA IN SMOKERS AND NONSMOKERS\*

Histologic Subtype	Smokers** (percent)	Non-smokers† (percent)
Squamous cell carcinoma	98.0	2.0
Adenocarcinoma	81.6	18.4
Bronchioloalveolar carcinoma	70.6	29.4
Small cell carcinoma	98.9	1.1
Large cell carcinoma	93.3	6.7

\*Modified from reference 9.

\*\*No. = 2708.

†No. = 218.

Some 80 percent of lung cancer deaths in men and 75 percent of lung cancer deaths in women can be attributed to cigarette smoking (38,45). In addition, it has been estimated that up to 25 percent of cases of carcinoma of the lung occurring in nonsmokers are the result of passive exposure to cigarette smoke (38), but this is controversial. In one study, 17 percent of the cases of lung carcinoma in nonsmokers were ascribed to childhood or adolescent exposure to cigarette smoke (31).

Asbestos is thought to be responsible for 4,000 to 6,000 deaths per year from carcinoma of the lung, less than 5 percent of all lung carcinoma deaths (38). A strong dose-response effect has been shown between asbestos exposure and the development of lung carcinoma; all histologic subtypes can be seen. There is a synergistic and multiplicative effect of smoking and asbestos exposure, with a 50-fold increased risk in the development of lung carcinoma (25). Although there is considerable debate, several studies suggest that the increased risk of lung carcinoma from asbestos exposure is only seen in those individuals with concomitant asbestosis (50).

Radiation is known to cause lung carcinoma (38). The association has been most extensively studied in uranium miners exposed to radon daughters. All types of lung carcinoma are seen, and a dose effect has been shown. An increased incidence of lung carcinoma has also been found in atom bomb survivors (35). Recently, considerable

2063632659

Table 7-5

**AGENTS OR EXPOSURES WITH PROVEN, PUTATIVE,  
OR POSSIBLE ASSOCIATION WITH LUNG CARCINOMA\***

Ionizing radiation	Tobacco smoke
Asbestos	Shipyards workers
Chloromethyl ether	Truck drivers
Cadmium	Plumbers
Arsenic	Rubber workers
Chromate	Coke oven workers
Hexavalent chromium	Petroleum workers
Formaldehyde	Mustard gas workers
Terpenes	Coal tar workers
Vinyl chloride	Roofers
Soots and tars	Pottery workers
Nickel	Printers
Isopropyl oils	Female cosmetologists
Antimony	Leather industry workers
Beryllium	Building laborers
Cobalt	Construction workers
Iron and iron oxides	Cooks, bakers, and pastry cooks
N-nitrosamines	Asbestos insulation workers
Polycyclic aromatic hydrocarbons	Uranium miners
Fibrous zeolites	
Manmade mineral fibers	Other factors:
Fiberglass	Lung scarring
Glass wool	Alveolar epithelial hyperplasia/ bronchioloalveolar cell adenoma
Rock wool	Previous bronchogenic carcinoma
Ceramics	Hereditary carcinomas (chapter 3)
Alumina	
Lead	

\*From references 17,25,28,35,38,39,42,43.

interest has centered on nonoccupational exposure to radon daughters in houses. Some studies have suggested that 5 to 15 percent of lung carcinomas are due to nonoccupational radon daughter exposure and that these cases account for 25 percent of the lung carcinomas occurring in nonsmokers and for 5 percent of the carcinomas in smokers (38). These figures have not been universally accepted (21), and further confirmatory studies are needed.

There has been some evidence that diet may have an impact on the risk of carcinoma of the lung. In a review by Colditz et al. (26), it was concluded that vitamin C might offer slight protection from lung carcinoma, but there was little evidence that vitamin C or vitamin E had any major influence. Data on a protective effect from selenium was limited but thought to warrant further study.

There are a number of other potential or proven agents or exposures associated with an increased risk of lung carcinoma, as shown in Table 7-5. In some instances, an occupation is associated with an increased incidence of carcinoma of the lung, but the specific causative agent or agents have not been identified. Some of the exposures listed in Table 7-5 are controversial and not universally accepted as proven or putative causes of lung carcinoma.

The concept of *scar carcinoma*, namely carcinoma of the lung arising in the vicinity of fibrosis (fig. 7-2), has been accepted for decades since first being studied in 1930s (17,36). In early studies, most of the scars were attributed to old tuberculosis; however, infarcts, pneumoconioses, and other chronic inflammatory scarring of the lung have all been implicated in individual



Figure 7-2  
SCAR CANCER

Small peripheral adenocarcinoma, bronchioloalveolar type, arising in association with a scar. There is a scar associated with some pleural puckering (arrows) around and from which the darker-appearing neoplasm extends. The surrounding lung tissue shows no significant fibrosis and only slight emphysema. In the majority of such cases, including this, the scar is probably the result of, rather than a precursor to, the carcinoma.

cases. In some studies, nearly half of all peripheral lung carcinomas were associated with a scar (36). The concept of scar carcinoma suggests that progressively atypical epithelial changes develop in the vicinity of the scar and ultimately lead to the development of a carcinoma. Meyer and Liebow (36) and subsequently others (27,51) described lung carcinomas arising in association with diffuse interstitial fibrosis and drew an analogy with carcinomas associated with focal scarring.

Tumors that have been labeled scar carcinomas are typically subpleural adenocarcinomas associated with retraction or puckering of the overlying pleura (17,22). Cut sections grossly show a central sclerotic zone ("scar") which may have anthracotic pigment and necrosis. Larger tumors (greater than 3 cm) tend to have more extensive scarring (22). In 1980, Shimosato (49) and subsequently others (22,32-34) suggested that the fibrosis in most scar carcinomas is a secondary phenomenon rather than a precursor to the carcinoma. Studies showed that the scar had abundant type III collagen and an extracellular matrix, suggesting an ongoing fibrosing process supporting a host response to the neoplasm (19,34), i.e., these tumors were desmoplastic carcinomas rather than carcinomas arising in scars.

Nevertheless, there are well-documented cases (52) showing carcinomas arising adjacent to old granulomas (fig. 7-3). It seems likely, however, that most scars are a secondary reaction rather than a preexisting lesion. This phenomenon is easily appreciated in sclerosing bronchioloalveolar carcinomas (chapter 13) in which only the alveolar walls involved by carcinoma are inflamed, thickened, and fibrosed.

While the pathogenesis of many scar carcinomas has been questioned, the concept of *atypical type 2 cell hyperplasia* (*atypical adenomatous hyperplasia*) as a precursor of adenocarcinoma has been reemphasized in several recent studies (23,40,41). Nakanishi (40) studied 15 cases of alveolar epithelial hyperplasia coexistent with lung carcinoma in a series of 70 patients. A histogenetic relationship between atypical alveolar epithelial hyperplasia (fig. 7-4) and pulmonary adenocarcinoma was suggested. Nakayama et al. (41) showed that atypical adenomatous hyperplasia (probably synonymous with *atypical alveolar epithelial hyperplasia*) was a clonal cellular proliferation closely related to well-differentiated adenocarcinoma.

In a series of 247 consecutive resections for carcinoma of the lung, Miller (37) found 23 (9.3

2063632661




Figure 7-3  
SCAR CANCER

Adenocarcinoma arising in association with a scar. These sections are from a physician's wife who had been followed for several decades with a stable lesion on her chest radiograph following a documented bout of tuberculosis. When the radiographic lesion started to enlarge, it was resected. Evidence of old healed granulomatous disease with increased anthracotic pigmentation can be seen at the left; in the surrounding fibrotic tissue, there is a proliferation of atypical epithelium (lower left and right) which was continuous with an obvious bronchioloalveolar carcinoma extending from the edge of the fibrotic region. This case probably represents a bona fide scar carcinoma.

percent) with incidental 1- to 7-mm nodules separate from the main tumor mass. These nodules were composed of localized proliferations of type 2 cells and were termed *bronchioloalveolar cell adenomas* (chapter 13). Miller concluded that these adenomas might be an early or premalignant phase of glandular neoplasia with the potential for progression to carcinoma. She also suggested that the finding might explain some examples of multicentricity among lung carcinomas. In individual cases, atypical type 2 cell hyperplasia and bronchioloalveolar cell adenoma may be difficult to distinguish and part of a disease spectrum.

Patients with a history of a resected lung carcinoma have an appreciable risk of a second lung carcinoma. The risk is in the range of 2 percent per year for patients with long-term survival

(over 5 years) (44). Long-term survivors of small cell carcinoma have a particularly high risk, estimated at 5.6 percent per person year (29).

Bejui-Thivolet et al. (20) have used in situ hybridization to show the presence of human papilloma virus (HPV) DNA in well-differentiated squamous cell carcinomas of the lung. They studied 33 carcinomas and 10 bronchial squamous metaplasias with probes for HPV subtypes 6, 11, 16, and 18. Of the 43 cases studied, 7 showed the presence of HPV DNA, including 1 case of squamous cell metaplasia and 6 cases of squamous cell carcinoma. Fourteen of the 43 lesions studied showed condylomatous changes, and 6 of the 7 cases in which HPV was detected by in situ hybridization came from among these cases. The authors suggested that HPV infection might be potentially oncogenic in the lung as it

2063632662



Figure 7-4

**ATYPICAL ADENOMATOUS HYPERPLASIA**

There is a relatively uniform proliferation of low cuboidal cells lining alveolar walls. Obvious cytologic features of malignancy are not present. Such atypical adenomatous hyperplasia is thought by some to represent a precursor to the development of adenocarcinoma of the lung, and such proliferations are relatively common in lungs resected for bronchioloalveolar carcinoma, as was this case.

is in the genital tract, but that its exact role needs to be clarified.

The morphogenesis of squamous cell carcinoma has been studied for several decades and is illustrated cytologically in figures 7-5 and 7-6. Since cigarette smoking increases squamous metaplasia (46), most studies have used serial sputum cytology specimens from cigarette smokers (16,18) or uranium miners. There is loss of normal ciliated lining cells with basal cell hyperplasia, low columnar nonciliated epithelium or squamous metaplasia, and increasing degrees of atypical squamous metaplasia (dysplasia), ultimately followed by squamous cell carcinoma in situ (CIS) and invasive squamous carcinoma. Not all cases of CIS of the bronchial tree are progressive, and some cases regress (24).

In practice, CIS is not found in all patients with squamous cell carcinoma and rarely in other histologic subtypes of lung carcinoma. In meticulous studies of serial sections of the bronchial tree taken at autopsy, Auerbach (18) found CIS in 26 of 34 patients with carcinoma of the lung. It is much less frequently seen in surgical specimens, as shown in the data of Rilke et al. (47): CIS at the bronchial margin was seen in only 3 of 67 cases (4.5 percent).

The molecular biology of carcinogenesis in the lung is discussed in chapter 3.

**SCREENING FOR CARCINOMA  
OF THE LUNG**

In 1971, the National Cancer Institute organized the Cooperative Early Lung Cancer Group to develop a screening program for the early detection of lung cancer in high-risk patients (male smokers) (53-59). Over 30,000 men older than 45 years of age enrolled in the study; at the time of entry into the study, none were suspected of having a lung carcinoma. Sputum cytologic examinations, chest radiographic examination, or both, along with a medical questionnaire were the screening tests used.

At the initial screening, a number of unsuspected lung carcinomas were identified, and these comprised the prevalence cancers in the study. The proportion of cases in each histologic subtype was roughly similar to that for all cases of lung carcinoma. Fifty-nine percent of the tumors were resectable, a figure more than double that seen in routine practice.

The carcinomas that were detected during the study period represented the incidence cancers. A number of both the prevalence cancers and the incidence cancers represented early or occult lung carcinomas, which are described and illustrated in chapter 10.

The study showed that with screening there was increased cancer detection, resectability, and survival. Regarding the incidence cancers: less than half were actually detected by the screening studies despite rescanning every 4 months; cancers detected on the basis of clinical symptoms were rarely resectable; and the proportion that could be completely resected was 46 percent for the screened group and 32 percent for the control group (58).

2063632663



Figure 7-5  
MORPHOGENESIS OF CARCINOMA OF THE LUNG .

Mild (A-C), moderate (D-F), and severe (G-I) squamous atypia (dysplasia) developing in uranium miners followed with sputum cytology examinations. (Courtesy of Drs. G. Saccomanno and M. Turner, Grand Junction, CO.)



Figure 7-5 (Continued)

2063632665





Figure 7-5 (Continued)

2063632666






Figure 7-6

MORPHOGENESIS OF BRONCHOGENIC CARCINOMA

Even though degrees of squamous atypia (dysplasia) may progress to squamous cell carcinoma (A-C), squamous atypia may also precede small cell carcinoma (D-F) or adenocarcinoma (G-I) in uranium miners. (Courtesy of Drs. G. Saccomanno and M. Turner, Grand Junction, CO.)

2063632667



Figure 7-6 (Continued)

2063632668



Figure 7-6 (Continued)

2063632669

Table 7-6

DETECTION OF LUNG CARCINOMA  
BY SCREENING METHODS\*

	% Detected By:		
	X Ray Only	Cytology Only	X Ray & Cytology
Squamous cell (no. = 81)	38	43	19
Adenocarcinoma (no. = 43)	81	0	19
Small cell (no. = 15)	73	0	27
Large cell (no. = 20)	75	10	15

\*From reference 54.

The screening methods showed considerable variability in effectiveness for identifying histologic subtypes of lung carcinoma (Table 7-6). Over 60 percent of the squamous cell carcinomas had a positive cytology, and in data not shown, it was found that 95 percent of all cancers detected by cytology alone were squamous cell carcinomas (54). For the other histologic subtypes, the chest radiograph was the primary means of detection.

The 5-year survival data from the Mayo Lung Project revealed the following (56): cases identified cytologically, about 80 percent; cases identified roentgenographically, about 35 percent; symptomatic cases, about 10 percent; every 4-month screened group, about 35 percent; and control group, less than 15 percent.

Despite these findings, there was no statistically significant difference in the mortality rate for lung carcinoma between the study groups and the control groups, either in the Mayo Lung Project or in the other two institutions taking part in the study (56). For the Mayo Lung Project there were 122 lung cancer deaths in the study group (3.2 per 1000 man years) and 115 lung cancer deaths in the control group (3.0 per 1000

man years). Although tumors were detected in the study group and overall there was a better 5-year survival, the fact that the mortality rates were similar between the study group and the control group suggests that some of the tumors identified in the study group may have been relatively indolent and slower growing and might not have manifested for a number of years or been fatal. It is possible that these tumors had inherent biologic differences from the more highly aggressive tumors that accounted for the mortality in both groups and characterize most lung carcinomas.

Other general conclusions from the Cooperative Early Lung Cancer Group study were (54): 1) the chest roentgenogram is the most sensitive method available for detecting lung carcinoma, and 2) sputum cytology is the most effective method for detecting early squamous cell carcinoma of the lung.

These studies suggest that mass screening does not have a significant impact on mortality, and thus question the usefulness of mass screening for carcinoma of the lung. This conclusion has recently been challenged by Strauss et al. (61) who concluded, after reworking the original data, that there was insufficient evidence to firmly recommend against lung cancer screening.

While mass screening may not have a significant impact on mortality, screening of select populations may yet prove to be of value. Saccomanno et al. (60) described the results of periodic cytologic examinations of sputum in a group of uranium miners, a subset now known to be at high risk for the development of lung carcinoma. In this group, abnormalities progressed in degree over time and changes were readily detectable in sputum cytology specimens (figs. 7-5, 7-6); there was an average of 4 to 5 years during which individuals would exfoliate abnormal cells prior to the development of an invasive carcinoma. The authors concluded that this 4- to 5-year period represented a window of opportunity for early detection and treatment.

2063632670

## REFERENCES

### Overview

1. Auerbach O, Garfinkel L. The changing pattern of lung carcinoma. *Cancer* 1991;68:1973-7.
2. Carey FA, Donnelly SC, Walker WS, Cameron EW, Lamb D. Synchronous primary lung cancers: prevalence in surgical material and clinical implications. *Thorax* 1993;48:344-6.
3. el-Torky M, el-Zeky F, Hall JC. Significant changes in the distribution of histologic types of lung cancer. A review of 4928 cases. *Cancer* 1990;65:2361-7.
4. Gazdar AF, Linnoila RI. The pathology of lung cancer—changing concepts and newer diagnostic techniques. *Semin Oncol* 1988;15:215-25.
5. Ichinose Y, Hara N, Ohta M. Synchronous lung cancers defined by deoxyribonucleic acid flow cytometry. *J Thorac Cardiovasc Surg* 1991;102:418-24.
6. Johnston WW. Histologic and cytologic patterns of lung cancer in 2580 men and women over a 15-year period. *Acta Cytol* 1988;32:163-8.
7. Linnoila I. Pathology of non-small cell lung cancer. New diagnostic approaches. *Hematol Oncol Clin North Am* 1990;4:1027-51.
8. Martini N, Melamed MR. Multiple primary lung cancers. *J Thorac Cardiovasc Surg* 1975;70:606-12.
9. Rohwedder JJ, Weatherbee L. Multiple primary bronchogenic carcinoma with a review of the literature. *Am Rev Respir Dis* 1974;109:435-45.
10. Rosenow EC, Carr DT. Bronchogenic carcinoma. *Cancer J Clin* 1979;29:233-45.
11. Teeter SM, Holmes FF, McFarlane MJ. Lung carcinoma in the elderly population. Influence of histology on the inverse relationship of stage to age. *Cancer* 1987;60:1331-6.
12. Travis WD, Travis LB, Percy C, Devesa SS. Lung cancer incidence and survival by histologic type. *Cancer*. In press.
13. Vincent RG, Pickren JW, Lane WW, et al. The changing histopathology of lung cancer: a review of 1682 cases. *Cancer* 1977;39:1647-55.

### Incidence

14. Boring CC, Squires TS, Tong T. Cancer statistics, 1993. *CA Cancer J Clin* 1993;43:7-26.
15. Garfinkel L, Silverberg E. Lung cancer and smoking trends in the United States over the past 25 years. *CA Cancer J Clin* 1991;41:137-45.

### Etiology

17. Auerbach O, Garfinkel L, Parks VR. Histologic type of lung cancer in relation to smoking habits, year of diagnosis, and sites of metastases. *Chest* 1975;67:382-7.
18. Garfinkel L, Parks VR. Scar cancer of the lung: increase over a 21-year period. *Cancer* 1979;43:636-42.
16. Gere JB, Forman JB, et al. Changes in the bronchial epithelium in relation to smoking and cancer of the lung. *N Engl J Med* 1957;256:97-104.
19. Barsky SH, Huang SJ, Bhuta S. The extracellular matrix of pulmonary scar carcinomas is suggestive of a desmoplastic origin. *Am J Pathol* 1986;124:412-9.
20. Bejui-Thivolet F, Liagre N, Chignol MC, Chardonnet Y, Patricot LM. Detection of human papilloma virus DNA in squamous bronchial metaplasia and squamous cell carcinomas of the lung by in situ hybridization using biotinylated probes in paraffin-embedded specimens. *Hum Pathol* 1990;12:111-6.
21. Bowie C, Bowie SH. Radon and health. *Lancet* 1991;337:409-13.
22. Cagle PT, Cohle SD, Greenberg SD. Natural history of pulmonary scar cancers. Clinical and pathologic implications. *Cancer* 1985;56:2031-5.
23. Carey FA, Wallace WA, Fergusson RJ, Kerr KM, Lamb D. Alveolar atypical adenomatous hyperplasia in association with primary pulmonary adenocarcinoma: a clinicopathologic study of ten cases. *Thorax* 1992;47:1041-3.
24. Carter D. Squamous cell carcinoma of the lung: an update. *Semin Diagn Pathol* 1985;2:226-34.
25. Churg A, Greene FH. Pathology of occupational lung disease. New York: Igaku-Shoin. 1988:50.
26. Colditz GA, Stampfer MJ, Willett WC. Diet and lung cancer. A review of the epidemiologic evidence in humans. *Arch Intern Med* 1987;147:157-60.
27. Fraire AE, Greenberg SD. Carcinoma and diffuse interstitial fibrosis of lung. *Cancer* 1973;31:1078-86.
28. Frank AL. The epidemiology and etiology of lung cancer. *Clin Chest Med* 1982;3:219-28.
29. Heyne KH, Lippman SM, Lee JJ, Lee JS, Hong WK. The incidence of second primary tumors in long-term survivors of small-cell lung cancer. *J Clin Oncol* 1992;10:1519-24.
30. Hoffmann D, Hoffmann I, Wynder EL. Lung cancer and the changing cigarette. *IARC Sci Publ* 1991;105:449-59.
31. Janerich DT, Thompson WD, Varela LR, et al. Lung cancer and exposure to tobacco smoke in the household. *N Engl J Med* 1990;323:632-6.
32. Kolin A, Koutoulakis T. Role of arterial occlusion in pulmonary scar cancers. *Hum Pathol* 1988;19:1161-7.
33. Kung IT, Lui IO, Loke SL, et al. Pulmonary scar cancer. The pathologic reappraisal. *Am J Surg Pathol* 1985;9:391-400.
34. Madri JA, Carter D. Scar cancers of the lung: origin and significance. *Hum Pathol* 1984;15:625-31.
35. Marchevsky AM. Pathogenesis and experimental models of lung cancer. *Lung Biol Health Dis* 1990;44:7-27.
36. Meyer EC, Liebow AA. Relationship of interstitial pneumonia honeycombing and atypical epithelial proliferation to cancer of the lung. *Cancer* 1965;18:322-51.
37. Miller RR. Bronchioloalveolar cell adenomas. *Am J Surg Pathol* 1990;14:904-12.
38. Minna JD, Pass H, Glatstein E, Ihde DC. Cancer of the lung. In: DeVita VT, Hellman S, Rosenberg SA, eds. *Cancer: principles and practice of oncology*. Philadelphia: JB Lippincott. 1988:591-705.
39. Mohsenifar Z. Epidemiology of lung cancer. *Lung Biol Health Dis* 1990;44:1-5.

2063632671

40. Nakanishi K. Alveolar epithelial hyperplasia and adenocarcinoma of the lung. *Arch Pathol Lab Med* 1990;114:363-8.
41. Nakayama H, Noguchi M, Tsuchiya R, Kodama T, Shimosato Y. Clonal growth of atypical adenomatous hyperplasia of the lung: cytofluorometric analysis of nuclear DNA content. *Mod Pathol* 1990;3:314-20.
42. Nemery B. Metal toxicity in the respiratory tract. *Eur Respir J* 1990;3:202-19.
43. Owens AH, Abeloff MD. Neoplasms of the lung. In: Calabresi P, Schein PS, Rosenberg SA, eds. *Medical oncology: basic principles and clinical management of cancer*. New York: MacMillan, 1985:715-57.
44. Pairolero PC, Williams DE, Bergstralh EJ, Piehler JM, Bernatz PE, Payne WS. Postsurgical stage I bronchogenic carcinoma: morbid implications of recurrent disease. *Ann Thorac Surg* 1984;38:331-8.
45. Parkin DM, Sasco AJ. Lung cancer: worldwide variation in occurrence and proportion attributable to tobacco use. *Lung Cancer (Elsevier)* 1993;9:1-16.
46. Peters EJ, Morice R, Benner SE, et al. Squamous metaplasia of the bronchial mucosa and its relationship to smoking. *Chest* 1993;103:1429-32.
47. Rilke F, Carbone A, Clemente C, Pilotti S. Surgical pathology of resectable lung cancer. In: Muggia F, Rozenzweig M, eds. *Lung cancer: progress in therapeutic research*. New York: Raven Press, 1979:129-42.
48. Saccomanno G, Archer VE, Auerbach O, Saunders RP, Brennan LM. Development of carcinoma of the lung as reflected in exfoliated cells. *Cancer* 1974;33:256-70.
49. Shimosato Y, Suzuki A, Hashimoto T, et al. Prognostic implications of fibrotic focus (scar) in small peripheral lung cancers. *Am J Surg Pathol* 1980;4:365-73.
50. Sluis-Cremer GK, Bezuidenhout BN. Relation between asbestosis and bronchial cancer in amphibole asbestos miners. *Br J Ind Med* 1989;46:537-40.
51. Turner-Warwick M, Lebowitz M, Burrows B, Johnson A. Cryptogenic fibrosing alveolitis and lung cancer. *Thorax* 1980;35:496-9.
52. Yoneda K. Scar carcinomas of the lung in a histoplasmosis endemic area. *Cancer* 1990;65:164-8.

#### Screening

53. Berlin NI, Buncher CR, Fontana RS, Frost JK, Melamed MR. The National Cancer Institute Cooperative Early Lung Cancer Detection Program. Results of the initial screen (prevalence). Early lung cancer detection: introduction. *Am Rev Respir Dis* 1984;130:545-9.
54. Early lung cancer detection: summary and conclusions. *Am Rev Respir Dis* 1984;130:565-70.
55. Flehinger BJ, Melamed MR, Zaman MB, Heelan RT, Perchick WB, Martini N. Early lung cancer detection: results of the initial (prevalence) radiologic and cytologic screening in the Memorial Sloan-Kettering study. *Am Rev Respir Dis* 1984;130:555-60.
56. Fontana RS. Screening for lung cancer. Recent experience in the United States. In: Hansen HH, ed. *Lung cancer: basic and clinical aspects*. Boston: Martinus Nijhoff, 1986:91-111.
57. \_\_\_\_\_, Sanderson DR, Taylor WF, et al. Early lung cancer detection. Results of the initial (prevalence) radiologic and cytologic screening in the Mayo Clinic study. *Am Rev Respir Dis* 1984;130:561-5.
58. \_\_\_\_\_, Sanderson DR, Woolner LB, et al. Screening for lung cancer. A critique of the Mayo Lung Project. *Cancer* 1991;67:1155-64.
59. Frost JK, Ball WC Jr, Levin ML, et al. Early lung cancer detection: results of the initial (prevalence) radiologic and cytologic screening the Johns Hopkins study. *Am Rev Respir Dis* 1984;130:549-54.
60. Saccomanno G, Archer VE, Auerbach O, Saunders RP, Brennan LM. Development of carcinoma of the lung as reflected in exfoliated cells. *Cancer* 1974;33:256-70.
61. Strauss GM, Gleason RE, Sugarbaker DJ. Screening for lung cancer re-examined. A reinterpretation of the Mayo Lung Project randomized trial on lung cancer screening. *Chest* 1993;103:337S-41S.

◆◆◆

2063632672